



The Application of Genomics in Agriculture

Yufei Yang^{1,*}

¹ Beijing Technology and Business University, Beijing 100048, China

Abstract

With the growing global population and the challenges of environmental change, genomics has a crucial role to play in improving agricultural yield and sustainability. This article reviews the application of genomics in crop, animal husbandry, fishery and forestry in the past five years. In crop production, genomics has facilitated the genetic improvement of major crops such as rice, wheat, and maize, which have improved crop nutritional value, yield, and resistance to abiotic stresses. In the livestock industry, genomics research in poultry and other livestock has driven breeding improvements and enhanced animal health and productivity. Fisheries use genomic resources for population management and conservation to achieve sustainable development. In addition, the application of genomics in forestry also shows great potential. Overall, the application of genomics not only improves agricultural productivity, but also helps to address the challenges of global food security and environmental protection.

Keywords: breeding, agriculture, genome.

1 Introduction

Genomics is an interdisciplinary biological discipline that involves the collective characterization and quantitative study of all genes in an organism and the comparative study of different genomes, rather than the structure and function of a single gene [2]. As a data-driven science, it mainly uses machine learning to capture the dependencies between data and deduce new biological hypotheses [1].

In the context of a growing global population, the urgent problem for agricultural development is: how to produce more food with limited resources, while having less impact on the environment [3]. The human population is expected to reach 10 billion by 2050, but crop yields appear to be leveling off or even declining due to climate change and limited supply of arable land, with a 60 percent increase needed to feed the world's 10 billion people. Therefore, improving agricultural productivity and sustainability is critical for the world. There is an urgent need for continuous scientific breakthroughs and technological innovations to ensure global food security in the future [4]. Genomics is of great significance in breeding, animal husbandry and sustainable agriculture. Genomics can make great contributions to plant crop improvement and animal breeding improvement by intercrossing with computer science and other disciplines. At the same time, genomics can also improve the adaptability of crops to the environment, respond to climate change, and ensure global food security by developing sustainable crops that can adapt to environmental



Academic Editor:

Bin Guo

Submitted: 23 January 2025

Accepted: 17 March 2025

Published: 28 March 2025

Vol. 2, No. 1, 2025.

10.62762/ASFP.2025.111574

*Corresponding author:

✉ Yufei Yang

2303020124@st.btbu.edu.cn

Citation

Yang, Y. (2025). The Application of Genomics in Agriculture. *Agricultural Science and Food Processing*, 2(1), 26–46.



© 2025 by the Author. Published by Institute of Central Computation and Knowledge. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>).




Figure 1. Graphical summary.

changes [5]. As shown in Figure 1, genomics technologies such as whole-genome sequencing and genome-wide association studies (GWAS) have enabled the identification of key genes and quantitative trait loci (QTLs) associated with agriculturally important traits, leading to the development of crops with enhanced nutritional value, yield, and stress resistance. These advances demonstrate the power of genomics in facilitating precision breeding and sustainable agriculture to address global food security challenges.

2 The application of genomics in crop farming

2.1 Crops for food

2.1.1 Rice(*Oryza sativa*)

Rice is an important cereal and model plant. Rice genomics research can be divided into three approaches: structural genomics, functional

genomics and quantitative genomics [2]. Due to the complexity of genomics research, two approaches are commonly used in combination. The combination of structural genomics and functional genomics can comprehensively analyze the composition, structure and function of genes. Yang et al. [11] used the rice seedling HuizhanNano as the experimental material and found that *Pi2*, *Pi2*, *Pi2*, and *Pi2* genes play a key role in the strong disease resistance of huizhan using nanopore long-read technology. Using functional molecular markers, Xuan et al. [8] found a significant correlation between the blast resistance gene *Pita* and blast resistance, and proposed to extend the lifespan of resistant varieties by designing reasonable distribution and rotation of blast resistance genes.

The key advantage of integrating structural and quantitative genomics lies in precisely analyzing complex trait-genome associations through genomic

physical structure and quantitative variation. Das et al. [9] has enriched the genomics of flower organs in rice crops by using high-throughput whole genome sequencing technology, pre-processing and mapping of reads, and variation analysis. Liu et al. [10] used 180 rice germplasms with large yield variations as experimental materials to perform association analysis of genes related to grain weight by target sequence capture sequencing, and found that TGW-HC4 was the best haplotype combination. By using genome resequencing technology, Yu et al. [3] identified a large number of genes and genomic regions or QTL controlling green traits, which enabled green super rice to be breeding and promoted.

The integration of functional and quantitative genomics enables systematic gene function analysis and precise quantification of genetic effects. Lee et al. [6] combined QTL mapping and GWAS analysis techniques to analyze heading date genes and rice nutrition-related genes, respectively, and showed that heading date genes had pleiotropic effects on grain, and accelerated heading and nutrient-rich rice breeding could be achieved by locating mutations and mutagenesis. The application of genomics has improved the nutritional content of rice [6, 7].

Beyond integrating these approaches, alternative avenues for investigation exist. By reviewing the origin and domestication history of African rice, Wambugu et al. [12] clarified that the progress of genomics could continuously change the previous understanding of rice domestication, so as to had a more accurate understanding of rice domestication process from the genomic level, which can eventually guide crop improvement. Ahmad [13] reviewed the use of several genomic and transcriptomic approaches to understand the ability of rice to withstand abiotic stress. Zhang et al. [14] provided the eRice database as an online research platform for the rice research community. To provide its epigenomics and genomics resources and will continue to enrich epigenetics data. In summary, genomics advances rice breeding through enhanced nutrition, accelerated heading, and improved blast resistance.

2.1.2 Wheat (*Triticum aestivum* L.)

Wheat is the world's largest cultivated and most widely distributed food crop [15]. Modern biological techniques have been widely used in wheat breeding [16]. Firstly, molecular marker technology has been widely used in wheat breeding. Kumar et al. [17] and Shewry et al. [18] used

molecular marker technology to predict genotype and improve nutrient content of Indian wheat. They introduced the trait by identifying QTLs for dietary fiber content and created wild and weed relatives of wheat. Babu et al. [19] performed a cis-genic approach by molecular marker-assisted selection (MAS) to identify multiple anti-rust genes simultaneously.

Secondly, the technology related to sequencing is more and more emerging. Kayess et al. [20] used next-generation genome sequencing to conduct extensive gene screening for heat-tolerant wheat varieties, and combined phenomics and genomics to develop new varieties with high temperature tolerance. Li et al. [16] proposed that the IWGSC RefSeq v2.0 (the second version of the wheat reference sequence) had a nearly complete wheat genome sequence, which could isolate and characterized the genes for agronomic traits important for wheat molecular breeding in the future. Following this, Wang et al. [21] focused on understanding the origin and evolutionary history of wheat, applying resistance gene enrichment sequencing and k-mer-based association mapping markers to clone disease-causing genes and accumulate genomic data. The evolutionary history of wheat was revealed by genome analysis of wild and domesticated wheat [22]. In 2023, Sun et al. [23] reviewed the extensive application of MAS, QTL mapping, and GWAS in genomics-assisted breeding (GAB) in breeding programs. Thereafter, Roychowdhury et al. [24] reviewed methods to improve key agronomics traits such as yield, disease resistance, drought tolerance, and nutritional quality of wheat through precise genetic modification. Subedi et al. [25] performed a translational genomics study of grain size regulatory genes in wheat using the rice candidate gene *OsGS3* through the candidate gene approach. Boehm et al. [26] focused on foreign introgression during wheat improvement, developing a genomics-based chromosome engineering pipeline to improve the efficacy and throughput of foreign introgression. Rhodes et al. [27] put forward the urgent need to use the global genomic surveillance system to continuously track and monitor wheat blast and prevent the global spread of rice blast. Finally, several important wheat genomics-related platforms are proposed, including WheatOmics [28], Online database of single grain wheat [29]. Sehgal et al. [30] developed WheatOmics, a multi-omics platform enabling wheat functional genomics research through gene identification, transcript analysis, regulatory mapping, gene networks, mutant/variant analysis,

optimized for non-bioinformaticians.

2.1.3 Corn (*Zea mays*)

To overcome the challenges and ensure the sustainable increase of maize yield, genomics-based breeding is required [31]. First, gene sequencing has made a major contribution to the elucidation of various mechanisms. By combining PacBio Sequel and Illumina HiSeq sequencing, Wang et al. [32] assembled a more continuous genome of *Bipolaris maydis*, providing a valuable resource for studying leaf dieback disease in southern maize and helping to elucidate the molecular mechanisms of its pathogenicity. After further high-throughput sequencing (NGS), Farooqi et al. [33] aimed at maize abiotic stress tolerance traits, combined with phenotypic analysis, and compared gene expression and function under extreme conditions to understand the molecular mechanism of abiotic stress in maize. QTLs for cold tolerance, heat stress, salt stress, and drought were also mapped. Subsequently, based on long-read sequencing technology, Liu et al. [34] used maize as a model to achieve gap-free genome assembly. This can eliminate uncertainty between genes and their regulators and understand the internal structure of repetitive domains, thus laying the foundation for epigenome analysis and ultimately creating seamless telomere assemblies of complex genomes in the future. Yang et al. [35] used maize HapMap and genomewide CRISPR mutation libraries to provide insight into population structural variation and the fact that a subset of transposable elements are transcriptionally active and can be induced under stress conditions, and developed an editing platform for high-throughput gene targeting in maize. This will revolutionize maize functional genomics research. Hu et al. [36] used genome assembly and population genome analysis to identify candidate genes associated with sweet maize traits. Dang et al. [37] has conducted GWAS and genomic prediction analyses for plant structural traits. It is concluded that the common origin of modern sweet corn in the United States was northern Mexico, and the plant configuration characters of sweet corn and waxy corn were understood.

For the GWAS technique, Waqas et al. [38] combined it with QTL mapping to explore important chromosomal regions and genes in maize under drought and heat stress. Farooqi et al. [33] reviewed the progress of maize multi-omics in abiotic stress resistance and breeding, highlighting the application and potential of genomics to metabolomics in stress resistance research. Using GWAS and co-expression analyses of

root traits, Wu et al. [39] revealed genetic differences between aboveground agrultural traits and RSA traits and provided an important candidate gene resource for future maize root genetic improvement. Talukder et al. [40] targeted breeding of waxy maize by recessive wx1 introgression using three superior self-bred maize cultivars HKI323, HKI1105, and HKI1128. This maize has a shorter reproductive cycle, is rich in amylopectin, and can be used as a wx1 gene donor in breeding programs. Muntean et al. [31] outlined the application of approaches such as cis-genetics and epigenetics to enable maize breeding to move from domestication to precision breeding through assisted selection processes or precise breeding programs, and enable the development of new maize crops with high yield and quality performance as quickly as possible. McMillen et al. [41] targeted maize in the African region and used an ecophysiological model to disentangle the relationship between genes and the environment by integrating genomic information with physiology.

Finally, Woodhouse et al. [42] proposed MaizeGDB, a genomic database that can manage pan-genomic data, so that researchers can easily understand the structural and functional differences between maize and its direct relatives. Recently, Zhou et al. [43] proposed a method to dwarf maize. Based on the genetic relationship between d024, indel-4 and ZmCYP90D1, they proposed to breed a new type of maize dwarf germplasm by crossing the corn-teosinte - Gramineum allopolyploid.

2.1.4 Soybean (*Glycine max*)

In recent years, significant progress has been made in soybean genomics, revealing the role of epigenetics in soybean regulation, polyploidy and diploidization, and the 3D structure of the soybean genome [44]. Sun et al. [45] reviewed various aspects of soybean genomics, including the application of 3D genomics, which was very extensive and comprehensive. More importantly, the important breeding trait of soybean is the relationship between oil yield and protein content, and many scientists have made great efforts in this regard. Liu et al. [46] used resequencing and GWAS to reveal key loci for soybean protein content and oil production. Kumar et al. [47] applied low-coverage resequencing, genome selection (GS), QTL mapping, and GWAS to seek ways to improve soybean seed oil and protein simultaneously. Kim et al. [48] identified candidate genes associated with protein, oil, and amino acid content in 203 wild soybean germplasm by GWAS and detected Single

Nucleotide Polymorphism(SNP) markers for protein and oil content at loci on new chromosomes 15 and 20. Jia et al. [49] used SLAF-seq data for genome-wide association analysis, combined with transcriptome expression data, to identify three potential candidate genes with excellent haplotypes, These included Glyma.03G186200, Glyma.09G099500 and Glyma.18G248900, which facilitated potential MAS and improve oil production. In addition, for other important agrologic traits, Liu et al. [50] conducted individual *de novo* genome assembly of 26 representative soybean samples, constructed genome maps, and conducted pan-genome analysis, which revealed many new genetic variations and helped to find the correlation between genetic variations and candidate genes. Du et al. [51] proposed that genomics has played an important role in soybean resistance to biological stresses of Staygreen syndrome, soybean Mosaic virus, soybean cystine, soybean common cutworm, *Phytophthora soei* root rot and stem rot using GWAS, linkage analysis and QTL analysis. Bhat et al. [52] introduced the application of GAB in soybean improvement. Through QTL analysis, high-throughput genotyping and high-throughput digital phenotyping (HTP) platforms, the utility of GAB in soybean has been greatly increased. By combining CRISPR-Cas9 and cutting-edge sequencing technologies such as RNA-seq and GWAS, Razzaq et al. [53] have made precise alterations of genetic information to greatly improve important traits such as yield, nutritional value, and stress resistance. Johnson et al. [54] reviewed the advances in genomics before and after soybean improvement in the last decade. GWAS and GS were proposed to improve the utilization efficiency of micronutrients in soybean and solve the problem of micronutrient deficiency in global soil.

Finally, the application of database is also indispensable. Razzaq et al. [53] also proposed the Soybean Genetics and Genomics Database (SoyBase) with extensive soybean genomics data. The database updates and new tools and features of SoyBase were detailed later by Brown et al. [55].

2.2 Cash crop

2.2.1 Peanut (*Arachis hypogaea*)

Peanut is an important food, oilseed and feed crop [56]. The development of sequencing technology has brought great progress to peanut genomics. Li et al. [57] used Chinese peanut as experimental material and obtained the first draft of genome sequence of

Bacteroides peanuttiae strain YY187 isolated from leaves by PacBio Sequel sequencing technology. Contribute to understanding the genetic diversity of fungal pathogens and further comparative genomics. Bhat et al. [56] performed whole-genome resequencing on 179 peanut samples to reveal the genome-wide structural and functional characteristics of SNP. In 2024, Raza et al. [58] proposed that GAB combined with NGS methods, high-throughput trait related markers, HTP, MAB, and GS can be used to design new varieties faster and help *de novo* domestication. In 2024, using long read sequencing and RNA sequencing, Xue et al. [59] introduced Amon2.0, a higher quality, more continuous, complete and accurate monticola genome assembly method, which was helpful for future molecular-assisted breeding. Ma et al. [60] used tetraploid peanut (*Arachis Tifrunner*) as experimental material to construct genomic Simple Sequence Repeats(SSRs) and combined with computer e-PCR analysis to enrich the genomic data of peanut SSRs. Tang et al. [61] used qRT-PCR analysis and CRISPR/Cas9 technology to identify a total of 19 AhFatB genes for improving peanut oil quality.

The impact of external stress on peanut development cannot be ignored. Huang et al. [62] conducted genomic research on peanut biological stress, and summarized the response of peanut to a variety of pests and microbial pathogens and the progress of omics. For example, Marker-Assisted Selection (MAS) and markassisted backcross have been successfully applied in peanut. O-methyltransferases (OMTs) play important roles in biological metabolism and stress tolerance. Cai et al. [63] conducted a genome-wide study of OMTs to provide a theoretical basis for understanding peanut development and stress response. For the Virginia peanut population, Newman et al. [64] used PCR allele competition expansion detection method, GS of high-quality markers and molecular marker assisted selection (MAS) methods to conduct a specific study on its genomic data. Wang et al. [65] reviewed that omics, especially genomic data, plays an important role in understanding the molecular basis of peanut seed development and also contributes to understanding the genetic and epigenetic mechanisms of key agronomic traits.

2.2.2 Cotton (*Gossypium*)

Cotton is an important fiber crop and model system in the world [66]. Firstly, Manivannan et al. [67] comprehensively reviewed the progress of cotton genomics in recent years, including the

introduction of functional genes in cotton organelle genome and methods to improve cotton fiber quality, etc. In addition, the application of NGS, GS, 3D genomics and cotton pan-genome in cotton was introduced. However, Nadeem et al. [68] introduced the application of sequencing technology to cotton leaf rolling virus resistance and construction of mutation libraries induced by EMS (a chemical mutagen in breeding) in cotton, respectively [69]. Comparative genomics, developed by sequencing technology, has also been widely used. Using four *Bacillus* strains as experimental materials, Liu et al. [70] performed comparative genomic analysis and found that *Bacillus velezensis* was more abundant in secondary metabolites, which could effectively control verticillium wilt in cotton. Cohen et al. [71] used genome-wide comparative techniques to identify several nematode resistance genes, which in turn could be introduced and stacked into several upland cotton lineages. A more comprehensive understanding can be obtained by combining various genomic technologies, such as QTL, GWAS, RNA interference (RNAi) and virus-induced gene silencing, and 3D genomics. Naoumkina et al. [72] review the application of these techniques to improve both cotton fiber yield and quality. Iram described the application in drought tolerance of cotton [73]. Wen et al. [74] revealed the origin of cotton species and applied to the study of candidate genes related to fiber development, Yang et al. [75] reviewed future plans for breeding "super cotton". Wang et al. [76] used Shixiya No.1 (SXY1), an Asian cotton plant, as experimental material to identify and regulate candidate genes, which promoted the progress of cotton functional genomics. Shen et al. [77] obtained genomic data of 890 upland cotton varieties through genome-wide selective scanning detection, and analyzed that Introgression is affected by both artificial selection and environmental factors, which contributed to the understanding of the mechanism of intraspecific penetration.

Finally, the establishment of a database is very important for the study of cotton's association with genomics technology. In 2020, Yang et al. [78] proposed that a global data exchange platform should be established. After that, Zhang et al. [79] introduced a Gossypian genome database and detailed its functions and usage, such as the ability to search for orthologous gene ids. Liu et al. [80] constructed Fingerprint Finder by short-read sequencing, indicating that GWAS and fingerprint sites complement each other, which can more

comprehensively understand the population genomics of cotton.

2.3 Vegetable crops

2.3.1 Cabbage (*Brassica oleracea*)

Chinese cabbage is a nutritious and economically significant vegetable crop [81]. Chinese cabbage can also be used as a model crop [82]. Ma et al. [82] proposed that high-density Chinese cabbage genetically mutated populations (CCGMPs) could promote the progress of functional genomics and use Chinese cabbage as a model to explore the function of leafy vegetable crops. Recent advances in the search for the cabbage genome, Lv et al. [83], based on High-throughput/resolution chromosome conformation capture (Hi-C) 's technology, assembled a high-quality draft cabbage genome. Xu et al. [84] discovered the importance of genomic SSRs markers for genetic linkage analysis, MAS, and genetic mapping by mining a large number of potentially variable SSNS through genome-wide bioinformatics surveys. As a kind of natural functional pigment, carotenoids are very nutritious. Cao et al. [81] used comparative genomics to study carotenoid biosynthesis genes, which laid a foundation for understanding the pathway of carotenoid biosynthesis and breeding cabbage with high carotenoid content. Yin et al. [85] used genome-wide data and bioinformatics methods to perform gene structure analysis, cis-acting element analysis, and collinearity analysis of the HD-Zip gene family, and found that the HD-Zip gene family has a regulatory role in carotenoid content and heat stress tolerance. In addition, for other agronomic shapes of cabbage, Ji et al. [86] used population comparative genomics, RNA-seq analysis, and qRT-PCR experiments to identify six genes that can be used for MAS and also understand the mechanism of heading traits. Wang et al. [87] combined multiple genomic data to analyze the glucosinolates (GSLs) biosynthetic genes, which helps to understand the synthesis mechanism of GSLs and prepare for the breeding of cabbage with high GSLs. Xing et al. [88] comprehensively analyzed the pseudo-response regulators (PRR) genes of cabbage cultivar 'BoZG17-1' through multiple sequence alignment PRR gene, gene structure analysis and cis-regulatory element analysis. It is beneficial to the cultivation of resistant varieties. Xu et al. [89] used genome sequencing, complete sequence alignment, genome structure comparison and phylogenetic analysis to study the cabbage chloroplast genome, which provided high-quality genetic breeding

information. Chen et al. [90] studied the genome of Ogura CMS cabbage by gene sequencing, collinearity analysis and PCR amplification, and based on this, a new non-breeding cabbage, Bel CMS cabbage, was constructed. It has the advantages of less exogenous cytoplasm and no dead buds at the beginning of flowering. Finally, databases can provide rich genomic data, and Li et al. [91] introduced the content and functions of a newly developed database, the non-heading Chinese cabbage database, which will be combined with bioinformatics to facilitate comparative genomic analysis in the future.

2.3.2 Carrot (*Brassica oleracea*)

Carrot is a nutrient-rich vegetable [92]. First, in order to improve the breeding of carrots, Macko-Podgorni et al. [93] used open-pollinated carrots as experimental materials and found that the expression of DcDCAF1 could regulate storage root development through GWAS. Liu et al. [95] improved hybridization of carrots using MAS, introgression, and third generation sequencing. Duran et al. [94] used sequencing genotyping and probe analysis to construct a genetic map of fertility restoration genes. Second, comparative genomics was also important. Zhao et al. [96] performed NGS and analysis of *Aspergillus rhizobium* isolate CBR2 (the causative agent of black rot of carrot) and *Alternaria dauci* (the causative agent of *Alternaria* leaf blight), respectively. Advances in comparative genomic analysis of carrots have been promoted. Zhe et al. [97] identified candidate genes for the formation of trichomes by resequencing and qRT-PCR analysis, which identified 12 genes associated with trichomes. Hao et al. [92] elucidated that DcNAC genes in carrot respond to abiotic stress by multiple sequence alignment and phylogenetic tree analysis. Finally, Rolling et al. [98] presented the content and functionality of the comprehensive database CarrotOmics, such as its ability to provide a visual carrot reference genome and drive genome-assisted selection.

2.4 Melon and fruit crops

2.4.1 Apple (*Malus pumila* Mill)

Apple is a fruit of economic importance [99]. Many attempts had been made to address abiotic stresses. Liu et al. [100] used the 'Baleng Crab' cultivar as the experimental material, and developed two markers, SNP182G and SNP11G/SNP761A, to increase the salt-alkali tolerance of apple rootstock through SNP marker analysis and accurate prediction of salt-alkali stress damage index. Yao et al. [101]

used "Golden Delicious" as experimental material to perform genome-wide identification and construct phylogenetic relationships for the HSP20 gene family, and finally identified several HSP20 genes as candidates for heat stress, which could contribute to the development of heat-tolerant apple varieties. Verma et al. [102] reviewed the role of multiple gene families in drought and salt tolerance in apple and introduced some miRNA candidates under investigation. In addition to abiotic stresses, pests and diseases are also a serious threat to apples. Anthrax is the causative agent of apple bitter rot. Khodadadi et al. [103] performed phylogenetic analysis and secondary metabolic gene cluster analysis by whole genome sequencing combined with bioinformatics software such as fastANI, which provided valuable genomic resources of anthrax and was beneficial to the disease management of apple stash. Carneiro et al. [104] completed the draft genome sequence of the pathogen *Colletotrichum chrysophilum* by sequencing the DNA of the *chrysophilum* strain M932 using a two-end sequencing system. It helps to develop varieties that can resist renal leaf spot disease and bitter rot disease. In order to select high-quality varieties, Shen et al. [105] proposed a new bioinformatics tool, bulked segregant analysis tool for out-of-crossing species and a genomics aided prediction model, they had used it to predict several QTLs and candidate genes for key agrologic traits in apple. Kumar et al. [106] used *Malus domestica* and wild *Malus* germplasm as experimental materials to study the genetic relationship of various nutrients in apple, including ascorbic acid, through apple 20 K SNP array genotyping, combined with GWAS and metabolite epitope analysis. This facilitates targeted breeding to improve different nutrients. Keller et al. [107] used apple REFPOP as experimental material and used FruitPhenoBox for gene phenotype analysis and SNP genotyping to study the genetic basis of apple shape and color, which helps to select the appearance of apple when breeding. Chen et al. [99] reviewed the previous research on red meat apples and proposed that MdMYB10 was an important research direction for the selection of high-quality red meat apples.

Finally, apple genomics data and research methods are also constantly enriching. For The 'Red Fuji' apple, Peng et al. [108] used HiFi long reads and Hi-C reads to analyze the high-quality haplotype 'Red Fuji' apple genome, which was a valuable resource for comparative genomics. In 2021, Minamikawa

et al. [109] used 184 apple breeding lines as experimental materials to show that combining the data of Infinium and the GRAS-Di genotype system, GWAS and genomic prediction (GP) can produce more accurate results. Then in 2024, Minamikawa et al. [110], using seven founders as experimental materials, through SNP genotyping chip system, GP and GWAS methods, the breeding value of the founder haplotype was revealed and the accuracy of these methods was evaluated.

2.4.2 Watermelon (*Citrullus lanatus*)

Watermelon is the third largest fruit crop in the world [113]. Understanding the genetic changes in watermelon domestication history helped modern watermelon breeding scientists to generate their seamless genomes. Using 27 watermelon and its relatives as experimental materials, and found that watermelon was derived from *Citrullus lanatus subsp. Cordophanus*, which was helpful for analyzing good shape and introducing it into breeding. Renner et al. [112] used Illumina sequencing and other technologies to perform chromosome level genome assembly of Kordofan melon and found a large number of genome structural variations (SV), which can provide insights into genetic changes in watermelon domestication history and contribute to modern watermelon breeding. Yang et al. [113] used *Citrullus lanatus* cultivars to study the genetic map of watermelon by Perfect SNP, Target SNP-seq technology and construction of phylogenetic trees, which helps to promote the diversity of watermelon varieties. It helps to promote the diversity of watermelon varieties. In order to breed superior varieties, Yang et al. [114] used two watermelon mother species "1061" and "812" as experimental materials, and found the structural and functional properties of CIPRX gene family, such as its involvement in watermelon skin lignin synthesis, through qRT-PCR analysis and combined with multiple bioinformatics analyses. Yang et al. [113] used bulk segregant analysis Pipelines and RNA-Seq methods to identify important QTLs that could contribute to the development of drought-tolerant watermelon cultivars. Maraga et al. [115] used the hybrid progeny of BIL-53 and IIHR-140-152 as experimental materials, and identified two potential candidate genes for watermelon epidermal traits through QTL mapping and mapping, candidate gene analysis and sequencing, and SNP identification, Named Cla97C09G175150 and Cla97C09G175170. Mashilo et al. [116], for *C. lanatus*. The comparative analysis of traits, genetic characteristics and genomes

was helpful for gene integration breeding.

Finally, some watermelon-related databases have gradually emerged. Deng et al. [111] used four watermelon germplasm as experimental materials to perform Genome *de novo* assembly and Oxford Nanopore Technology long-read sequencing. High-quality watermelon reference genomes and mutation libraries were obtained. Patel et al. [117] summarized the history and mechanism of watermelon anthracnose and proposed that for sustainable cultivation of watermelon anthracnose resistance, it is necessary to establish resistance gene libraries to continuously improve resistance. The application of genomics in other crops is also summarized in Table 1.

3 The application of genomics to animal husbandry

3.1 Poultry

Zhang et al. [129] introduced the current status of genomic mating in poultry, which was a new method of animal breeding and still need continue to be combined with bioinformatics to handle large amounts of data in the future. Chicken (*Gallus gallus domesticus*) is the most widely farmed poultry. Li et al. [130] demonstrated that combining GWAS/SNPs and machine learning improved the accuracy of chicken genome prediction. Wang et al. [131] explored the field of RNA editing in chickens by establishing an RNA database of chickens combined with whole-genome analysis. In addition to these advances in genomic technologies, blight was also an important threat to chickens. Zhou et al. [132] introduced a genomics laboratory, GIP IL, which carried out genomic prediction and selection through GWAS and sequencing SNP panel. Excellent varieties resistant to Newcastle disease (ND) were bred. Smith [133] reviewed the application of genomics to animal defense against various avian viruses, such as ND, avian influenza, and poxvirus. Chen et al. [127] identified gene silencing by RNAi as a novel tool for mite control by analyzing core RNAi pathway genes in *Dermanyssus gallinae*. Feng et al. [134] used poultry in the southern United States as experimental materials to analyze *E. coli* through molecular markers and comparative genomics, which was helpful for the development of corresponding vaccines. In addition, in order to breed excellent varieties. Olaniyan et al. [135] reviewed methods to integrate community-based breeding programmes with genomics to facilitate breeding improvement in

Table 1. Other studies of genomics in crop farming.

Subjects	Technology	Conclusion	References
Lentils	NGS, MAS, SNP	Genomics for the promotion of lentil breeding, yield improvement, and adaptation to adverse environments.	Kumar et al. [118]
	miRNAomics, epigenomics, phenomics	Developing Saline-Smart Lentils.	Ali et al. [119]
	CRISPR/Cas9, QTL genotyping, GWAS, SNP	Review of drought tolerance mechanisms in lentils. The source of key genes responsible for salt tolerance in lentils was identified.	Saini et al. [120] Dissanayake et al. [121]
	disease resistance breeding, hybridization, mutagenesis, molecular markers	This review summarized recent advances in improving disease resistance and climate adaptation of lentils.	Roy et al. [122]
weeds	Genome Sequencing	Integrating multi-omics data can provide a comprehensive understanding of weed resistance mechanisms.	Patterson et al. [123]
	Molecular marker techniques, multi-omics techniques	To better understand the mechanisms of invasive weeds.	Huang et al. [124]
	The WeedPedia platform	This review summarized the current status of weed genomics and introduced the International Weed Genomics Consortium (IWGC).	Montgomery et al. [125]
	QTL, Genome assembly	The author aimed at cruciferous weeds to study the early-growth of weeds and understand the origin of weeds.	Li et al. [126]
Tomato	Repetitive sequence analysis, Genome assembly	The genomic basis for the enhanced herbicide adaptability of polyploid <i>Leptochloa chinensis</i> was explored.	Chen et al. [127]
	CRISPR/Cas9	Promoting plant research and precision plant breeding.	Xia et al. [128]

chickens. Wadood et al. [136] review the application of genomics to improved chicken reproduction, including techniques for epigenomics and precision gene editing. Long [137] investigated ways to improve the accuracy of predicting male fertility using genomic tools and identified markers such as the *TSPY* and *PRM* genes. Nematbakhsh et al. [138] designed a new breed of local village chicken with discrimination by genome-wide SNP. To prevent commercial fraud by selling colored broiler chickens as if they were local village chicken. High-quality databases are essential for genomics research. Barros et al. [139] used long-read technologies to identify new target genes and assemble a new turkey genome, providing a wealth of genomic data. Li et al. [140] presented for the first time a genomics database of helmeted guinea fowl and introduced its use. Li et al. [140] combined multiple sequencing techniques for *de novo* genome assembly and established a *de novo* pan-genome of 20 chickens to provide genomic data for breeding improvement of chickens.

Ducks (*Anas*) are also farmed in large numbers around the world. Che et al. [141] demonstrated the advantages of HiFi reads technology in identifying duck genomes by RNA-seq and identification of differentially expressed genes using healthy female mule ducks as experimental material. Guo et al. [142] explored the complex origin of domestic ducks by introgression analysis of whole genome sequences of mallards, maple leaf ducks, and domestic ducks. Chang et al. [143] used Chinese crested duck as

experimental material, and through construction genome, comparative genomics analysis and SV, adaptive evolution and the origin of CC duck were further explored. Lin et al. [144] used Chaohu and Ji'an Red ducks as experimental material to understand the differentiation process of ducks based on the analysis of population structure and candidate genes. To select for superior traits, Du et al. [145] combined genomics data with bioinformatics tools to investigate the molecular mechanisms by which ducks stop laying eggs, as well as changes in the liver that contribute to increased egg production. Zhang et al. [146] identified MC1R as the key gene regulating feather color through genotyping, SNP, and GWAS in a large number of ducks of different species. Yu et al. [147] investigated the mechanism of artificial selection by resequencing using Peking duck as experimental material.

Recent studies by various research teams have utilized genomic data and analyses to investigate the origin, introgression, and superior traits of different goose (*Anser cygnoides*) breeds, providing valuable insights and candidate genes for breeding improvements. Wen et al. [148] used the sequencing data of swan and greylag geese to understand the origin and introgression of Chinese geese through demographic analysis and candidate gene annotation analysis. Zhang et al. [149] performed a high-quality genome assembly of wild swan goose, providing a high-quality reference genome. Zhou et al. [150] targeted feather sac development in The Hungarian White Goose and generated high-quality genome-wide

data that could help improve breeding. Liu et al. [151] identified FAP and GCG as candidate genes for high egg production by signal analysis based on SNP and qRT-PCR, which simultaneously contributed to the selection of Wulong geese with high egg production. Zhang et al. [152] conducted Linkage disequilibrium analysis on genome-wide data and identified several candidate genes, such as TGFBR3L, that were associated with superior traits in geese, which could help goose breed improvement.

3.2 Domestic animals

Animal husbandry is important for the nutrient supply of the human diet, and the development of genomics is important for animal husbandry. Cows (*bovis*) can provide nutritious meat and milk. Guinan et al. [153] confirmed the role of GS on breeding improvement by analyzing genetic data from the National Cooperator Database and investigating multiple dairy cow breeds. Kertz et al. [154] theoretically reviewed the application of SNPS and other genomic methods to Improve Female Fertility, which still needs further practice in the future. Du et al. [155] reviewed the application of genomics in understanding the genetic mechanisms of cattle for main bovine species. Tian et al. [156] reviewed the role of GWAS, GP, and GS in breeding high-quality beef. Aiming at the quality and yield of dairy cattle, Zhao et al. [157] constructed the genome of predicted genes for cows with A2A2 casein genotype by CDHIT (v4.5.7) software, and analyzed their rumen fluids. The signature metabolite AA was found to play a key role in milk quality. Billa et al. [158] performed feed restriction by grouping experiments and extracted mirnas for microarray analyses and bioinformatic analyses, respectively, mirna was found to regulate milk yield by affecting mammary structure and lipid biosynthesis. Zavadilova et al. [159] used genome-wide association and genetic evaluation to improve disease resistance in dairy cows.

Pigs (*Sus scrofa*) are also one of the most important livestock. Yin et al. [160] provided valuable genomic data on Ningxiang pigs by performing Runs of homozygosity as well as genomic breed composition. Li et al. [161] performed gene introgression scan on Yunnan indigenous pigs and European pigs, through specific gene loci such as *NAF1*. The discovery of population gene flow in pigs helps to understand the genetic mechanisms behind traits and ultimately drives. Piorkowska et al. [162] review recent advances in pig genomics, including exploration of the origins of pig domestication, genes involved in lipogenesis,

and candidate genes for litter size. Johansson et al. [163] compared two types of porcine kidney epithelial samples by short-read sequencing and found aneuploidies, which contributed to further understanding of livestock genomics.

In addition, the horse (*Equus caballus*) industry is also an important part of the livestock industry. Flack et al. [164] used Oxford Nanopore sequencing to construct a high-quality genome assembly for The Przewalski's horse breed, providing valuable genomic data. Littiere et al. [165] identified 53 candidate genes through a genome-wide association study and candidate gene analysis to understand the genetic map of horses of various breeds. Polak et al. [166] demonstrated the application of genomics in exploring inbreeding in rare horses by pedigree analysis of Sokolski horses and Sztumski horses using biological software Admixture. Frohlich et al. [167] conducted a study on Timor ponies (TP) to investigate the genetic relationship of TP through genotyping, SNP and ROH detection, which is helpful for breeding studies of TP.

Finally, Saleem et al. [168] comprehensively reviewed the major development of Nutrigenomics in animal husbandry, which has broad development prospects for improving animal yield and traits by exploring the relationship between genes and nutrition. A summary of the application of genomics in other animals is provided in Table 2.

4 Applications of genomics to fisheries

Fishing is an important part of agriculture, with a rich diversity of fish species, but it also needs to be protected. To enrich fishery genomics resources, Alvarenga et al. [177] constructed 70 fish mitochondrial genomes through custom baits and target enrichment, which are molecular resources for fish management. Papa et al. [178] provided a high-quality population genomics resource by assembling the genome of the marine teleost tarakihi by combining short Illumina reads and long nanopore reads.

Bertram et al. [179] conducted SNP genotyping and bioinformatics analysis on snapper (*Lutjanus erythropterus*) from the coast of Western Australia, and investigated the population structure of snapper using population genomics. A more continuous genome assembly by long-read sequencing by Blommaert et al. [180] identified multiple structural variants, which is using genomics to facilitate fisheries management. Timm et al. [181] used community

Table 2. Other studies of genomics in to animal husbandry.

Subjects	Technology	Conclusion	References
Sheep,goat	GS, SNP, GWAS	Efficiency of genetic improvement to increase fertility in bulls.	Taylor et al. [169]
	Gene families clustering analysis, RNA sequencing	Understanding the taste of goat milk at the genomic level.	Zhang et al. [170]
	GWAS, QTP,	To understand the genetic mechanisms of adaptation to extreme conditions in goats.	Li et al. [171]
	WGS data, phylogenetic analysis	To explore gene flow through Y chromosome variation in goats.	Nijman et al. [172]
	Pedigree data, Phenotypic data	Comparison of breeding value and genomic differences between Sanin and Alpine goats.	Negro et al. [173]
	SNPs, detecting InDels	A goat genome-wide data platform, the VarGoats project, was introduced.	Denoyelle et al. [174]
chicken	single-nuclei ATAC-seq	This paper summarizes the achievements of single-cell omics research and analyzes the development prospects of this technology.	Lyons et al. [175]
	CRISPR/Cas9, base editing	To generate valuable poultry genome editing models.	Park et al. [176]

genomics to evaluate several oceanography models, including features such as spawning grounds, to help increase reproduction of Mexican Pink Shrimp (*Penaeus duorarum*) as well as to understand their evolutionary history. Ferrette et al. [182] performed *de novo* Genome Assembly and established a new reference genome for ssailfish (*Istiophorus albicans*), which is beneficial for understanding ssailfish speciation and may advance comparative genomic. Woodings et al. [183] used ddRADSeq and SNP to investigate the population structure of Eastern Rock Lobster to assess the harm of overfishing. Timm et al. [184] review the current issues to be addressed by Seascape genomics and its importance to maintaining a sustainable fishery. Oppen et al. [185] reviewed the current widespread use of genomics in marine conservation, demonstrating the need to use genomic monitoring and reference genomes to conserve marine diversity. Caccavo et al. [186] generated whole-genome libraries by small-scale sequencing of otoliths to provide genomic data for fisheries genomics. Casas et al. [187, 188] review the application of multiple emerging genomic technologies such as high throughput molecular technologies and RAD-seq in fisheries. Andersson et al. [189] reviewed methods of population genomics to promote sustainable fisheries, such as SNP-chip analysis and genome prediction, among others. Using golden perch as experimental material, Attard et al. [190] proposed that detection of ancestry of broodstock through genomics could promote the integration of genomics and fishery management. A summary of the application of genomics in other animals is provided in Table 2.

5 Applications of genomics to forestry

Forestry plays an important role both in protecting diversity and in the economy [191]. But forestry breeding has a longer cycle than crop breeding, so the application of genomics is even more important. Genomic technologies are widely used in forestry [192]. Cao et al. [191] reviewed the application of CRISPR-Cas in forest trees, which has been used in fruit trees but still needs technological breakthroughs to improve efficiency and quality.

Mostert et al. [193] studies on the domestication of wood fiber crops by SNP genotyping can provide insights into evolutionary history and population structure. Anders et al. [194] reviewed the role of Cas-DNA-free technology and multiplex editing in tree breeding improvement. Ahmar et al. [195] reviewed the role of genotyping and TILLING screening in GAB for advancing forest tree breeding. In addition, many scientists have provided valuable forest genomics resources. Burley et al. [196] studied two tree species *Jacaranda copaia* and *Handroanthus*. By combining Illumina short-read and Chicago libraries data and using whole genome alignment technology, we found gene duplication between them. Niu et al. [197] reviewed recent advances in poplar genomics, from haploid sequencing to the realization of telomere to telomere (T2T), through length long sequencing technology, helping to advance breeding. Tumas et al. [198] used SNP technology to construct genetic maps of *Picea sitchensis* and *Picea glauca*, which is helpful for the development of genotyping chips and sustainable management of forest tree species. Capblancq et al. [199] revealed the genetic history of *oriental beech* and provided a quality genomics model by studying its whole genome data in the Caucasus region. Luo et al. [200] reviewed *Dipterocarpaceae*

genomic data, combined with comparative genomics and phylogenetic analyses, to help select superior *Dipterocarpaceae* tree species using GS. Han et al. [201] established and analyzed the genome structure of *Quercus variabilis* by PacBio HiFi sequencing, which is conducive to comparative genomics studies. Matallana-Ramirez et al. [202] used Loblolly Pine as an example to illustrate the important role of genomics in managing forest ecosystems by discussing the development status of phenotypical genotype studies. Shirasawa et al. [203] analyzed four conifer species by HiFi long-read sequencing and found associations among them, helping to speed up the breeding process through genomic data. Finally, the impact of climate change on forests is also significant. Based on high-throughput phenotyping technology, Feng et al. [204] proposed combining landscape genomics and evolutionary genomics to cultivate trees that intelligently respond to climate change. Sharma et al. [192] reviewed the application of genotyping, QTL, and genome prediction in tree breeding. Abiri et al. [205] reviewed the effects of rainfall and other environmental stresses on forest trees, as well as the analysis of tolerance mechanisms of forest trees.

6 Conclusion

In the past five years, the application of genomics in agriculture has achieved remarkable results, especially in improving crop yield, quality, stress tolerance and adaptability, and promoting sustainable development of animal husbandry and fisheries. With the help of GAB technology, scientists can accurately improve crop and animal varieties to effectively meet the dual challenges of global food demand growth and environmental protection.

In the future, the application of genomics in agriculture will be more promising. With the continuous progress of technology and the emergence of new tools, genomics is expected to further enhance agricultural productivity, improve the quality and safety of agricultural products, and provide strong support for the sustainable development of agriculture and the environment. The integration of genomics with emerging technologies such as big data, artificial intelligence and biotechnology will inject new impetus into agricultural innovation and transformation. Continued investment in genomics research and its application in agriculture is important to meet global food demand, ensure food security and protect the environment. In addition, the potential of genomics to address climate change, such as breeding drought and

heat tolerant crop varieties and protecting biodiversity through optimized population management and habitat restoration, deserves further exploration. The future of agriculture will rely on genomics to build more resilient, productive, and sustainable food systems for present and future generations.

Data Availability Statement

Not applicable.

Funding

This work was supported without any funding.

Conflicts of Interest

The author declare no conflicts of interest.

Ethical Approval and Consent to Participate

Not applicable.

References

- [1] Eraslan, G., Avsec, Ž., Gagneur, J., & Theis, F. J. (2019). Deep learning: new computational modelling techniques for genomics. *Nature reviews genetics*, 20(7), 389-403. [CrossRef]
- [2] Wang, C., & Han, B. (2022). Twenty years of rice genomics research: From sequencing and functional genomics to quantitative genomics. *Molecular Plant*, 15(4), 593-619. [CrossRef]
- [3] Yu, S., Ali, J., Zhou, S., Ren, G., Xie, H., Xu, J., ... & Zhang, Q. (2022). From Green Super Rice to green agriculture: Reaping the promise of functional genomics research. *Molecular Plant*, 15(1), 9-26. [CrossRef]
- [4] Gao, C. (2021). Genome engineering for crop improvement and future agriculture. *Cell*, 184(6), 1621-1635. [CrossRef]
- [5] Naglaa A, A. (2022). Genome editing techniques in plants: a comprehensive review and future prospects toward zero hunger.
- [6] Lee, S. Y., Jeung, J. U., & Mo, Y. (2024). Allelic combinations of Hd1, Hd16, and Ghd7 exhibit pleiotropic effects on agronomic traits in rice. *G3: Genes, Genomes, Genetics*, 14(3), jkad300. [CrossRef]
- [7] Zaghumi, M. J., Ali, K., & Teng, S. (2022). Integrated genetic and omics approaches for the regulation of nutritional activities in rice (*Oryza sativa* L.). *Agriculture*, 12(11), 1757. [CrossRef]
- [8] Xuan, N., Zhang, H., Liu, X., Guo, Q., Liu, J., & Yao, F. (2020). Analysis of the relationship between blast resistance genes and disease resistance of rice germplasm via functional molecular markers. *Phyton*, 89(1), 45. [CrossRef]

- [9] Das, S. P., Jasrotia, R. S., Deb, D., Iquebal, M. A., Jaiswal, S., & Dey, N. (2021). Genomic analysis of polycarpellary rice (*Oryza sativa* L.) through whole genome resequencing. *Journal of Plant Biochemistry and Biotechnology*, 30, 364-372. [CrossRef]
- [10] Liu, G., Qiu, D., Lu, Y., Wu, Y., Han, X., Jiao, Y., ... & Zhang, Z. (2023). Identification of Superior Haplotypes and Haplotype Combinations for Grain Size and Weight-Related Genes for Breeding Applications in Rice (*Oryza sativa* L.). *Genes*, 14(12), 2201. [CrossRef]
- [11] Yang, W., Yang, Z., Yang, L., Li, Z., Zhang, Z., Wei, T., ... & Li, G. (2024). Genomic and transcriptomic analyses of the elite rice variety Huizhan provide insight into disease resistance and heat tolerance. *Genomics*, 116(5), 110915. [CrossRef]
- [12] Wambugu, P. W., Ndjiondjop, M. N., & Henry, R. (2021). Genetics and genomics of African rice (*Oryza glaberrima* Steud) domestication. *Rice*, 14(1), 6. [CrossRef]
- [13] Ahmad, M. (2022). Genomics and transcriptomics to protect rice (*Oryza sativa* L.) from abiotic stressors: pathways to achieving zero hunger. *Frontiers in Plant Science*, 13, 1002596. [CrossRef]
- [14] Zhang, P., Wang, Y., Chachar, S., Tian, J., & Gu, X. (2020). eRice: a refined epigenomic platform for japonica and indica rice. *Plant Biotechnology Journal*, 18(8), 1642. [CrossRef]
- [15] Jin, L., Li, Z., & Zhang, J. (2024). Research on Plant Genomics and Breeding 2.0. *International Journal of Molecular Sciences*, 25(12), 6659.
- [16] Li, J., Yang, J., Li, Y., & Ma, L. (2020). Current strategies and advances in wheat biology. *The Crop Journal*, 8(6), 879-891. [CrossRef]
- [17] Kumar, S., Jacob, S. R., Mir, R. R., Vikas, V. K., Kulwal, P., Chandra, T., ... & Singh, K. (2022). Indian Wheat genomics initiative for harnessing the potential of Wheat germplasm resources for breeding disease-resistant, nutrient-dense, and climate-resilient cultivars. *Frontiers in genetics*, 13, 834366. [CrossRef]
- [18] Shewry, P. R., Lovegrove, A., Wingen, L. U., & Griffiths, S. (2022). Opinion Exploiting genomics to improve the benefits of wheat: Prospects and limitations. *Journal of Cereal Science*, 105, 103444. [CrossRef]
- [19] Babu, P., Baranwal, D. K., Harikrishna, Pal, D., Bharti, H., Joshi, P., ... & Singh, A. (2020). Application of genomics tools in wheat breeding to attain durable rust resistance. *Frontiers in Plant Science*, 11, 567147. [CrossRef]
- [20] Kayess, O., Ashrafuzzaman, M., Khan, M. A. R., & Siddiqui, M. N. (2024). Functional phenomics and genomics: Unravelling heat stress responses in wheat. *Plant Stress*, 100601. [CrossRef]
- [21] Wang, Z., Miao, L., Chen, Y., Peng, H., Ni, Z., Sun, Q., & Guo, W. (2023). Deciphering the evolution and complexity of wheat germplasm from a genomic perspective. *Journal of genetics and genomics*, 50(11), 846-860. [CrossRef]
- [22] Ahmed, H. I., Heuberger, M., Schoen, A., Koo, D. H., Quiroz-Chavez, J., Adhikari, L., ... & Krattinger, S. G. (2023). Einkorn genomics sheds light on history of the oldest domesticated wheat. *Nature*, 620(7975), 830-838. [CrossRef]
- [23] Sun, C., Hu, H., Cheng, Y., Yang, X., Qiao, Q., Wang, C., ... & Chen, F. (2023). Genomics-assisted breeding: the next-generation wheat breeding era. *Plant Breeding*, 142(3), 259-268. [CrossRef]
- [24] Roychowdhury, R., Ghatak, A., Kumar, M., Samantara, K., Weckwerth, W., & Chaturvedi, P. (2024). Accelerating wheat improvement through trait characterization: advances and perspectives. *Physiologia Plantarum*, 176(5), e14544. [CrossRef]
- [25] Subedi, M., Ghimire, B., Bagwell, J. W., Buck, J. W., & Mergoum, M. (2023). Wheat end-use quality: State of art, genetics, genomics-assisted improvement, future challenges, and opportunities. *Frontiers in Genetics*, 13, 1032601. [CrossRef]
- [26] Boehm Jr, J., & Cai, X. (2024). Enrichment and diversification of the wheat genome via alien introgression. *Plants*, 13(3), 339. [CrossRef]
- [27] Rhodes, J. (2023). Genomic surveillance urgently needed to control wheat blast pandemic spreading across continents. *Plos Biology*, 21(4), e3002090. [CrossRef]
- [28] Ma, S., Wang, M., Wu, J., Guo, W., Chen, Y., Li, G., ... & Ni, F. (2021). WheatOmics: A platform combining multiple omics data to accelerate functional genomics studies in wheat. *Molecular plant*, 14(12), 1965-1968. [CrossRef]
- [29] Sharma, P. K., Ahmed, H. I., Heuberger, M., Koo, D. H., Quiroz-Chavez, J., Adhikari, L., ... & Tiwari, V. K. (2023). An online database for einkorn wheat to aid in gene discovery and functional genomics studies. *Database*, 2023, baad079. [CrossRef]
- [30] Sehgal, D., Dhakate, P., Ambreen, H., Shaik, K. H. B., Rathan, N. D., Anusha, N. M., ... & Vikram, P. (2023). Wheat omics: advancements and opportunities. *Plants*, 12(3), 426. [CrossRef]
- [31] Muntean, L., Ona, A., Berindean, I., Racz, I., & Muntean, S. (2022). Maize breeding: from domestication to genomic tools. *Agronomy*, 12(10), 2365. [CrossRef]
- [32] Wang, Y., Kang, H., Yao, J., Li, Z., Xia, X., Zhou, S., & Liu, W. (2022). An improved genome sequence resource of *Bipolaris maydis*, causal agent of Southern Corn Leaf Blight. *Phytopathology*, 112(6), 1386-1390. [CrossRef]
- [33] Farooqi, M. Q. U., Nawaz, G., Wani, S. H., Choudhary, J. R., Rana, M., Sah, R. P., ... & Siddique, K. H. (2022). Recent developments in multi-omics and breeding strategies for abiotic stress tolerance in maize (*Zea mays* L.). *Frontiers in plant science*, 13, 965878.

- [CrossRef]
- [34] Liu, J., Seetharam, A. S., Chougule, K., Ou, S., Swentowsky, K. W., Gent, J. I., ... & Dawe, R. K. (2020). Gapless assembly of maize chromosomes using long-read technologies. *Genome Biology*, 21, 1-17. [CrossRef]
- [35] Yang, N., & Yan, J. (2021). New genomic approaches for enhancing maize genetic improvement. *Current Opinion in Plant Biology*, 60, 101977. [CrossRef]
- [36] Hu, Y., Colantonio, V., Müller, B. S., Leach, K. A., Nanni, A., Finegan, C., ... & Resende Jr, M. F. (2021). Genome assembly and population genomic analysis provide insights into the evolution of modern sweet corn. *Nature Communications*, 12(1), 1227. [CrossRef]
- [37] Dang, D., Guan, Y., Zheng, H., Zhang, X., Zhang, A., Wang, H., ... & Qin, L. (2023). Genome-wide association study and genomic prediction on plant architecture traits in sweet corn and waxy corn. *plants*, 12(2), 303. [CrossRef]
- [38] Waqas, M. A., Wang, X., Zafar, S. A., Noor, M. A., Hussain, H. A., Azher Nawaz, M., & Farooq, M. (2021). Thermal stresses in maize: effects and management strategies. *Plants*, 10(2), 293. [CrossRef]
- [39] Wu, B., Ren, W., Zhao, L., Li, Q., Sun, J., Chen, F., & Pan, Q. (2022). Genome-wide association study of root system architecture in maize. *Genes*, 13(2), 181. [CrossRef]
- [40] Talukder, Z. A., Muthusamy, V., Chhabra, R., Bhatt, V., Reddappa, S. B., Mishra, S. J., ... & Hossain, F. (2022). Enrichment of amylopectin in sub-tropically adapted maize hybrids through genomics-assisted introgression of waxy1 gene encoding granule-bound starch synthase (GBSS). *Journal of Cereal Science*, 105, 103443. [CrossRef]
- [41] McMillen, M. S., Mahama, A. A., Sibiya, J., Lübberstedt, T., & Suza, W. P. (2022). Improving drought tolerance in maize: Tools and techniques. *Frontiers in Genetics*, 13, 1001001. [CrossRef]
- [42] Woodhouse, M. R., Cannon, E. K., Portwood, J. L., Harper, L. C., Gardiner, J. M., Schaeffer, M. L., & Andorf, C. M. (2021). A pan-genomic approach to genome databases using maize as a model system. *BMC plant biology*, 21, 1-10. [CrossRef]
- [43] Zhou, Y., Li, Y., Luo, L., Zhang, D., Wang, X., Chen, Y., ... & Tang, Q. (2025). Maize-Tripsacum-Teosinte allopolyploid (MTP), a novel dwarf mutant inducer tool in maize. *Plant Biotechnology Journal*, 23(1), 112-127. [CrossRef]
- [44] Zenda, T., Liu, S., Dong, A., & Duan, H. (2021). Advances in cereal crop genomics for resilience under climate change. *Life*, 11(6), 502. [CrossRef]
- [45] Sun, Z., Lam, H. M., Lee, S. H., Li, X., & Kong, F. (2024). Soybean functional genomics: bridging theory and application. *Molecular Breeding*, 44(1), 2. [CrossRef]
- [46] Liu, S., Liu, Z., Hou, X., & Li, X. (2023). Genetic mapping and functional genomics of soybean seed protein. *Molecular Breeding*, 43(4), 29. [CrossRef]
- [47] Kumar, V., Vats, S., Kumawat, S., Bisht, A., Bhatt, V., Shivaraj, S. M., ... & Sonah, H. (2021). Omics advances and integrative approaches for the simultaneous improvement of seed oil and protein content in soybean (*Glycine max* L.). *Critical Reviews in Plant Sciences*, 40(5), 398-421. [CrossRef]
- [48] Kim, W. J., Kang, B. H., Kang, S., Shin, S., Chowdhury, S., Jeong, S. C., ... & Ha, B. K. (2023). A genome-wide association study of protein, oil, and amino acid content in wild soybean (*Glycine soja*). *Plants*, 12(8), 1665. [CrossRef]
- [49] Jia, H., Han, D., Yan, X., Zhang, L., Liang, J., & Lu, W. (2024). Genome-wide association and RNA-Seq analyses reveal a potential candidate gene related to oil content in soybean seeds. *International Journal of Molecular Sciences*, 25(15), 8134. [CrossRef]
- [50] Liu, Y., Du, H., Li, P., Shen, Y., Peng, H., Liu, S., ... & Tian, Z. (2020). Pan-genome of wild and cultivated soybeans. *Cell*, 182(1), 162-176. [CrossRef]
- [51] Du, H., Fang, C., Li, Y., Kong, F., & Liu, B. (2023). Understandings and future challenges in soybean functional genomics and molecular breeding. *Journal of Integrative Plant Biology*, 65(2), 468-495. [CrossRef]
- [52] Bhat, J. A., & Yu, D. (2021). High-throughput NGS-based genotyping and phenotyping: Role in genomics-assisted breeding for soybean improvement. *Legume Science*, 3(3), e81. [CrossRef]
- [53] Razzaq, M. K., Babur, M. N., Awan, M. J. A., Raza, G., Mobeen, M., Aslam, A., & Siddique, K. H. (2024). Revolutionizing soybean genomics: How CRISPR and advanced sequencing are unlocking new potential. *Functional & Integrative Genomics*, 24(5), 153. [CrossRef]
- [54] Johnson, K. H., Krishna, T. A., Dash, M., Thiyageshwari, S., Ceasar, S. A., & Selvi, D. (2023). Food and nutritional security: innovative approaches for improving micronutrient use efficiency in soybean (*Glycine max* (L.) Merrill) under hostile soils. *Journal of Soil Science and Plant Nutrition*, 23(1), 56-70. [CrossRef]
- [55] Brown, A. V., Conners, S. I., Huang, W., Wilkey, A. P., Grant, D., Weeks, N. T., ... & Nelson, R. T. (2021). A new decade and new data at SoyBase, the USDA-ARS soybean genetics and genomics database. *Nucleic Acids Research*, 49(D1), D1496-D1501. [CrossRef]
- [56] Bhat, R. S., Shirasawa, K., & Chavadi, S. D. (2022). Genome-wide structural and functional features of single nucleotide polymorphisms revealed from the whole genome resequencing of 179 accessions of *Arachis*. *Physiologia Plantarum*, 174(1), e13623. [CrossRef]
- [57] Li, S., Xue, X., Gao, M., Wang, N., Cui, X., Sang, S., ... & Wang, Z. (2021). Genome resource for peanut web blotch causal agent *Peyronellaea arachidicola* strain

- YY187. *Plant Disease*, 105(4), 1177-1178. [CrossRef]
- [58] Raza, A., Chen, H., Zhang, C., Zhuang, Y., Sharif, Y., Cai, T., ... & Zhuang, W. (2024). Designing future peanut: the power of genomics-assisted breeding. *Theoretical and Applied Genetics*, 137(3), 66. [CrossRef]
- [59] Xue, H., Zhao, K., Zhao, K., Han, S., Chitkineni, A., Zhang, L., ... & Yin, D. (2024). A near complete genome of *Arachis monticola*, an allotetraploid wild peanut. *Plant biotechnology journal*, 22(8), 2110. [CrossRef]
- [60] Ma, J., Zhao, Y., Chen, H., Fu, C., Zhu, L., Zhou, X., ... & Zhao, C. (2020). Genome-wide development of polymorphic microsatellite markers and their application in peanut breeding program. *Electronic Journal of Biotechnology*, 44, 25-32. [CrossRef]
- [61] Tang, Y., Huang, J., Ji, H., Pan, L., Hu, C., Qiu, X., ... & Qiao, L. (2022). Identification of AhFatB genes through genome-wide analysis and knockout of AhFatB reduces the content of saturated fatty acids in peanut (*Arachis hypogaea* L.). *Plant Science*, 319, 111247. [CrossRef]
- [62] Huang, R., Li, H., Gao, C., Yu, W., & Zhang, S. (2023). Advances in omics research on peanut response to biotic stresses. *Frontiers in plant science*, 14, 1101994. [CrossRef]
- [63] Cai, T., Sharif, Y., Zhuang, Y., Yang, Q., Chen, X., Chen, K., ... & Zhuang, W. (2023). In-silico identification and characterization of O-methyltransferase gene family in peanut (*Arachis hypogaea* L.) reveals their putative roles in development and stress tolerance. *Frontiers in Plant Science*, 14, 1145624. [CrossRef]
- [64] Newman, C. S., Andres, R. J., Youngblood, R. C., Campbell, J. D., Simpson, S. A., Cannon, S. B., ... & Dunne, J. C. (2023). Initiation of genomics-assisted breeding in Virginia-type peanuts through the generation of a de novo reference genome and informative markers. *Frontiers in Plant Science*, 13, 1073542. [CrossRef]
- [65] Wang, Z., Lei, Y., & Liao, B. (2024). Omics-driven advances in the understanding of regulatory landscape of peanut seed development. *Frontiers in Plant Science*, 15, 1393438. [CrossRef]
- [66] Zhang, T., Wang, S., Chen, J., Zhu, S., Zhu, Q., & Zhao, T. (2024). Advances in Cotton Genomics, Genetics and Breeding. *Plants*, 13(18), 2579. [CrossRef]
- [67] Manivannan, A., & Cheeran Amal, T. (2023). Deciphering the complex cotton genome for improving fiber traits and abiotic stress resilience in sustainable agriculture. *Molecular Biology Reports*, 50(8), 6937-6953. [CrossRef]
- [68] Nadeem, S., Riaz Ahmed, S., Luqman, T., Tan, D. K., Maryum, Z., Akhtar, K. P., ... & Liu, Y. (2024). A comprehensive review on *Gossypium hirsutum* resistance against cotton leaf curl virus. *Frontiers in Genetics*, 15, 1306469. [CrossRef]
- [69] Wang, P., Abbas, M., He, J., Zhou, L., Cheng, H., & Guo, H. (2024). Advances in genome sequencing and artificially induced mutation provides new avenues for cotton breeding. *Frontiers in Plant Science*, 15, 1400201. [CrossRef]
- [70] Liu, H., Zeng, Q., Yalimaimaiti, N., Wang, W., Zhang, R., & Yao, J. (2021). Comprehensive genomic analysis of *Bacillus velezensis* AL7 reveals its biocontrol potential against *Verticillium* wilt of cotton. *Molecular Genetics and Genomics*, 296, 1287-1298. [CrossRef]
- [71] Cohen, Z. P., Perkin, L. C., Wagner, T. A., Liu, J., Bell, A. A., Arick, M. A., ... & Suh, C. P. (2024). Nematode-resistance loci in upland cotton genomes are associated with structural differences. *G3: Genes, Genomes, Genetics*, 14(9), jkae140. [CrossRef]
- [72] Naoumkina, M., & Kim, H. J. (2023). Bridging molecular genetics and genomics for cotton fiber quality improvement. *Crop Science*, 63(4), 1794-1815. [CrossRef]
- [73] Sharif, I., Aleem, S., Junaid, J. A., Ali, Z., Aleem, M., Shahzad, R., ... & Ellahi, F. (2024). Multiomics approaches to explore drought tolerance in cotton. *Journal of Cotton Research*, 7(1), 32. [CrossRef]
- [74] Wen, X., Chen, Z., Yang, Z., Wang, M., Jin, S., Wang, G., ... & Zhu, Y. (2023). A comprehensive overview of cotton genomics, biotechnology and molecular biological studies. *Science China Life Sciences*, 66(10), 2214-2256. [CrossRef]
- [75] Yang, Z., Gao, C., Zhang, Y., Yan, Q., Hu, W., Yang, L., ... & Li, F. (2023). Recent progression and future perspectives in cotton genomic breeding. *Journal of Integrative Plant Biology*, 65(2), 548-569. [CrossRef]
- [76] Wang, W., Yang, X., Zeng, J., Liang, A., Liu, D., Wang, W., ... & Xiao, Y. (2024). Construction of ethyl methane sulfonate mutant library in *G. arboreum* and rapid identification of mutant genes via repeated re-sequencing. *Industrial Crops and Products*, 213, 118373. [CrossRef]
- [77] Shen, C., Cao, Z., Xu, Z., Ouyang, L., Zhang, X., Guo, Z., ... & Huang, W. (2023). Population genomics unravels the characteristic relationship between introgression and geographical distribution in upland cotton. *Agronomy*, 13(7), 1781. [CrossRef]
- [78] Yang, Z., Qanmber, G., Wang, Z., Yang, Z., & Li, F. (2020). *Gossypium* genomics: trends, scope, and utilization for cotton improvement. *Trends in Plant Science*, 25(5), 488-500. [CrossRef]
- [79] Zhang, Z., Chai, M., Yang, Z., Yang, Z., & Fan, L. (2022). GRAND: an integrated genome, transcriptome resources, and gene network database for *gossypium*. *Frontiers in Plant Science*, 13, 773107. [CrossRef]
- [80] Liu, S., Cheng, H., Zhang, Y., He, M., Zuo, D., Wang, Q., ... & Song, G. (2024). Fingerprint Finder: Identifying Genomic Fingerprint Sites in Cotton Cohorts for Genetic Analysis and Breeding Advancement. *Genes*, 15(3), 378. [CrossRef]
- [81] Cao, W., Wang, P., Yang, L., Fang, Z., Zhang, Y., Zhuang, M., ... & Ji, J. (2021). Carotenoid biosynthetic

- genes in cabbage: Genome-wide identification, evolution, and expression analysis. *Genes*, 12(12), 2027. [CrossRef]
- [82] Ma, W., Zhang, P., Zhao, J., & Hong, Y. (2023). Chinese cabbage: an emerging model for functional genomics in leafy vegetable crops. *Trends in Plant Science*, 28(5), 515-518. [CrossRef]
- [83] Lv, H., Wang, Y., Han, F., Ji, J., Fang, Z., Zhuang, M., ... & Yang, L. (2020). A high-quality reference genome for cabbage obtained with SMRT reveals novel genomic features and evolutionary characteristics. *Scientific reports*, 10(1), 12394. [CrossRef]
- [84] Xu, Y., Xing, M., Song, L., Yan, J., Lu, W., & Zeng, A. (2021). Genome-wide analysis of simple sequence repeats in cabbage (*Brassica oleracea* L.). *Frontiers in plant science*, 12, 726084. [CrossRef]
- [85] Yin, L., Sun, Y., Chen, X., Liu, J., Feng, K., Luo, D., ... & Zhao, J. (2023). Genome-wide analysis of the hd-zip gene family in Chinese cabbage (*Brassica rapa* subsp. *pekinensis*) and the expression pattern at high temperatures and in carotenoids regulation. *Agronomy*, 13(5), 1324. [CrossRef]
- [86] Ji, G., Long, Y., Cai, G., Yan, G., Wu, J., Zhang, F., ... & Wu, X. (2023). Whole-genome comparison reveals structural variations behind heading leaf trait in *Brassica oleracea*. *International Journal of Molecular Sciences*, 24(4), 4063. [CrossRef]
- [87] Wang, P., Cao, W., Yang, L., Zhang, Y., Fang, Z., Zhuang, M., ... & Ji, J. (2023). Glucosinolate biosynthetic genes of cabbage: Genome-wide identification, evolution, and expression analysis. *Genes*, 14(2), 476. [CrossRef]
- [88] Xing, Y., Jiang, Y., Muhammad, A. R., & Song, J. (2024). Genome-wide identification of pseudo-response regulator (PRR) family members in cabbage (*Brassica oleracea* var. *capitata* L.) and their expression in response to abiotic stress. *The Journal of Horticultural Science and Biotechnology*, 99(2), 168-178. [CrossRef]
- [89] Xu, Y., Xing, M., Li, J., Zeng, A., Song, L., & Yan, J. (2022). Complete chloroplast genome sequence and variation analysis of *Brassica oleracea* L. *Acta Physiologiae Plantarum*, 44(10), 106. [CrossRef]
- [90] Chen, L., Ren, W., Zhang, B., Chen, W., Fang, Z., Yang, L., ... & Zhang, Y. (2021). Organelle comparative genome analysis reveals novel alloplasmic male sterility with orf112 in *Brassica oleracea* L. *International Journal of Molecular Sciences*, 22(24), 13230. [CrossRef]
- [91] Li, Z., Li, Y., Liu, T., Zhang, C., Xiao, D., & Hou, X. (2022). Non-heading Chinese cabbage database: An open-access platform for the genomics of *Brassica campestris* (syn. *Brassica rapa*) ssp. *chinensis*. *Plants*, 11(8), 1005. [CrossRef]
- [92] Hao, J. N., Wang, Y. H., Duan, A. Q., Liu, J. X., Feng, K., & Xiong, A. S. (2020). NAC family transcription factors in carrot: genomic and transcriptomic analysis and responses to abiotic stresses. *DNA and Cell Biology*, 39(5), 816-827. [CrossRef]
- [93] Macko-Podgórn, A., Stelmach, K., Kwolek, K., Machaj, G., Ellison, S., Senalik, D. A., ... & Grzebelus, D. (2020). Mining for candidate genes controlling secondary growth of the carrot storage root. *International Journal of Molecular Sciences*, 21(12), 4263. [CrossRef]
- [94] Duran, S. T., & Ipek, A. (2022). Identification of SNP markers linked to Rf locus in carrot using GBS. *Turkish Journal of Agriculture and Forestry*, 46(6), 898-907. [CrossRef]
- [95] Liu, X., Zhao, D., Ou, C., Hao, W., Zhang, Y., He, Y., ... & Zhuang, F. (2023). Draft genome sequence of carrot Alternaria leaf blight pathogen *Alternaria dauci*. *Plant Disease*, 107(7), 2197-2200. [CrossRef]
- [96] Zhao, D., Zhuang, F., Ou, C., Zhang, Y., & Liu, X. (2023). Complete genome sequence of the carrot black rot pathogen *Alternaria radicina* isolate CBR2. *Microbial Pathogenesis*, 185, 106421. [CrossRef]
- [97] Zhe, W. U., Xuan, Y. A. N. G., Yu-xuan, Z. H. A. O., & Li, J. I. A. (2022). Identifying candidate genes involved in trichome formation on carrot stems by transcriptome profiling and resequencing. *Journal of Integrative Agriculture*, 21(12), 3589-3599. [CrossRef]
- [98] Rolling, W. R., Senalik, D., Iorizzo, M., Ellison, S., Van Deynze, A., & Simon, P. W. (2022). CarrotOmics: A genetics and comparative genomics database for carrot (*Daucus carota*). *Database*, 2022, baac079. [CrossRef]
- [99] Chen, Z., Yu, L., Liu, W., Zhang, J., Wang, N., & Chen, X. (2021). Research progress of fruit color development in apple (*Malus domestica* Borkh.). *Plant Physiology and Biochemistry*, 162, 267-279. [CrossRef]
- [100] Liu, J., Shen, F., Xiao, Y., Fang, H., Qiu, C., Li, W., ... & Han, Z. (2020). Genomics-assisted prediction of salt and alkali tolerances and functional marker development in apple rootstocks. *BMC genomics*, 21, 1-16. [CrossRef]
- [101] Yao, F., Song, C., Wang, H., Song, S., Jiao, J., Wang, M., ... & Bai, T. (2020). Genome-wide characterization of the HSP20 gene family identifies potential members involved in temperature stress response in apple. *Frontiers in Genetics*, 11, 609184. [CrossRef]
- [102] Verma, S., Dubey, N., Mishra, V., Kumar, S., Sharma, R., Sharma, S., ... & Thakur, A. K. (2024). Functional Genomics of Salt and Drought Stress Tolerance in the Temperate Crop Apple (*Malus domestica*). *Journal of Plant Growth Regulation*, 43(11), 3941-3957. [CrossRef]
- [103] Khodadadi, F., Luciano-Rosario, D., Gottschalk, C., Jurick, W. M., & Ćimović, S. G. (2024). Unveiling the Arsenal of Apple Bitter Rot Fungi: Comparative Genomics Identifies Candidate Effectors, CAZymes, and Biosynthetic Gene Clusters in *Colletotrichum* Species. *Journal of Fungi*, 10(7), 493. [CrossRef]
- [104] Amaral Carneiro, G., Cali, M., Cappelletti, E., Baroncelli, R., & Prodi, A. (2023). Draft genome sequence of the apple pathogen *Colletotrichum chrysophilum* strain M932. *Journal of Plant Pathology*,

- 105(3), 1141-1143. [CrossRef]
- [105] Shen, F., Bianco, L., Wu, B., Tian, Z., Wang, Y., Wu, T., ... & Zhang, X. (2022). A bulked segregant analysis tool for out-crossing species (BSATOS) and QTL-based genomics-assisted prediction of complex traits in apple. *Journal of Advanced Research*, 42, 149-162. [CrossRef]
- [106] Kumar, S., Molloy, C., Hunt, M., Deng, C. H., Wiedow, C., Andre, C., ... & McGhie, T. (2022). GWAS provides new insights into the genetic mechanisms of phytochemicals production and red skin colour in apple. *Horticulture Research*, 9, uhac218. [CrossRef]
- [107] Keller, B., Jung, M., Bühlmann-Schütz, S., Hodel, M., Studer, B., Broggini, G. A., & Patocchi, A. (2024). The genetic basis of apple shape and size unraveled by digital phenotyping. *G3: Genes, Genomes, Genetics*, 14(5), jkae045. [CrossRef]
- [108] Peng, H., Yi, Y., Li, J., Qing, Y., Zhai, X., Deng, Y., ... & Zheng, Y. (2024). A haplotype-resolved genome assembly of *Malus domestica* 'Red Fuji'. *Scientific Data*, 11(1), 592. [CrossRef]
- [109] Minamikawa, M. F., Kunihiya, M., Moriya, S., Shimizu, T., Inamori, M., & Iwata, H. (2024). Genomic prediction and genome-wide association study using combined genotypic data from different genotyping systems: application to apple fruit quality traits. *Horticulture Research*, 11(7), uhae131. [CrossRef]
- [110] Minamikawa, M. F., Kunihiya, M., Noshita, K., Moriya, S., Abe, K., Hayashi, T., ... & Iwata, H. (2021). Tracing founder haplotypes of Japanese apple varieties: application in genomic prediction and genome-wide association study. *Horticulture Research*, 8. [CrossRef]
- [111] Deng, Y., Liu, S., Zhang, Y., Tan, J., Li, X., Chu, X., ... & Zhang, X. (2022). A telomere-to-telomere gap-free reference genome of watermelon and its mutation library provide important resources for gene discovery and breeding. *Molecular plant*, 15(8), 1268-1284. [CrossRef]
- [112] Renner, S. S., Wu, S., Pérez-Escobar, O. A., Silber, M. V., Fei, Z., & Chomicki, G. (2021). A chromosome-level genome of a Kordofan melon illuminates the origin of domesticated watermelons. *Proceedings of the National Academy of Sciences*, 118(23), e2101486118. [CrossRef]
- [113] Yang, J., Zhang, J., Du, H., Zhao, H., Mao, A., Zhang, X., ... & Xu, Y. (2022). Genetic relationship and pedigree of Chinese watermelon varieties based on diversity of perfect SNPs. *Horticultural Plant Journal*, 8(4), 489-498. [CrossRef]
- [114] Yang, T., Zhang, P., Pan, J., Amanullah, S., Luan, F., Han, W., ... & Wang, X. (2022). Genome-wide analysis of the peroxidase gene family and verification of lignin synthesis-related genes in watermelon. *International Journal of Molecular Sciences*, 23(2), 642. [CrossRef]
- [115] Maragal, S., Nagesh, G. C., Reddy, D. L., & Rao, E. S. (2022). QTL mapping identifies novel loci and putative candidate genes for rind traits in watermelon. *3 Biotech*, 12(2), 46. [CrossRef]
- [116] Mashilo, J., Shimelis, H., Maja, D., & Ngwepe, R. M. (2023). Meta-analysis of qualitative and quantitative trait variation in sweet watermelon and citron watermelon genetic resources. *Genetic Resources and Crop Evolution*, 70(1), 13-35. [CrossRef]
- [117] Patel, T., Quesada-Ocampo, L. M., Wehner, T. C., Bhatta, B. P., Correa, E., & Malla, S. (2023). Recent Advances and Challenges in Management of *Colletotrichum orbiculare*, the Causal Agent of Watermelon Anthracnose. *Horticultrae*, 9(10), 1132. [CrossRef]
- [118] Kumar, S., Rajendran, K., Kumar, J., Hamwieh, A., & Baum, M. (2015). Current knowledge in lentil genomics and its application for crop improvement. *Frontiers in plant science*, 6, 78. [CrossRef]
- [119] Ali, F., Zhao, Y., Ali, A., Waseem, M., Arif, M. A., Shah, O. U., ... & Wang, Z. (2024). Omics-driven strategies for developing saline-smart lentils: a comprehensive review. *International Journal of Molecular Sciences*, 25(21), 11360. [CrossRef]
- [120] Saini, S., Sharma, P., Sharma, J., Pooja, P., & Sharma, A. (2024). Drought stress in *Lens culinaris*: effects, tolerance mechanism, and its smart reprogramming by using modern biotechnological approaches. *Physiology and Molecular Biology of Plants*, 30(2), 227-247. [CrossRef]
- [121] Dissanayake, R., Cogan, N. O., Smith, K. F., & Kaur, S. (2021). Application of genomics to understand salt tolerance in lentil. *Genes*, 12(3), 332. [CrossRef]
- [122] Roy, A., Sahu, P. K., Das, C., Bhattacharyya, S., Raina, A., & Mondal, S. (2023). Conventional and new-breeding technologies for improving disease resistance in lentil (*Lens culinaris* Medik). *Frontiers in Plant Science*, 13, 1001682. [CrossRef]
- [123] Patterson, E. L., Saski, C., Küpper, A., Beffa, R., & Gaines, T. A. (2019). Omics potential in herbicide-resistant weed management. *Plants*, 8(12), 607. [CrossRef]
- [124] Huang, P., Hameed, R., Abbas, M., Balooch, S., Alharthi, B., Du, Y., ... & Du, D. (2023). Integrated omic techniques and their genomic features for invasive weeds. *Functional & Integrative Genomics*, 23(1), 44. [CrossRef]
- [125] Montgomery, J., Morran, S., MacGregor, D. R., McElroy, J. S., Neve, P., Neto, C., ... & Gaines, T. A. (2024). Current status of community resources and priorities for weed genomics research. *Genome biology*, 25(1), 139. [CrossRef]
- [126] Li, L. Z., Xu, Z. G., Chang, T. G., Wang, L., Kang, H., Zhai, D., ... & Wang, J. W. (2023). Common evolutionary trajectory of short life-cycle in Brassicaceae ruderal weeds. *Nature Communications*, 14(1), 290. [CrossRef]
- [127] Chen, K., Yang, H., Peng, Y., Liu, D., Zhang, J., Zhao,

- Z., ... & Wang, L. (2023). Genomic analyses provide insights into the polyploidization-driven herbicide adaptation in *Leptochloa* weeds. *Plant Biotechnology Journal*, 21(8), 1642-1658. [CrossRef]
- [128] Xia, X., Cheng, X., Li, R., Yao, J., Li, Z., & Cheng, Y. (2021). Advances in application of genome editing in tomato and recent development of genome editing technology. *Theoretical and Applied Genetics*, 134(9), 2727-2747. [CrossRef]
- [129] Zhang, P., Qiu, X., Wang, L., & Zhao, F. (2022). Progress in genomic mating in domestic animals. *Animals*, 12(18), 2306. [CrossRef]
- [130] Li, X., Chen, X., Wang, Q., Yang, N., & Sun, C. (2024). Integrating bioinformatics and machine learning for genomic prediction in chickens. *Genes*, 15(6), 690. [CrossRef]
- [131] Wang, Y. M., Ye, L. Q., Wang, M. S., Zhang, J. J., Khederzadeh, S., Irwin, D. M., ... & Wu, D. D. (2022). Unveiling the functional and evolutionary landscape of RNA editing in chicken using genomics and transcriptomics. *Zoological research*, 43(6), 1011. [CrossRef]
- [132] Zhou, H., Baltenweck, I., Dekkers, J., Gallardo, R., Kayang, B. B., Kelly, T., ... & Lamont, S. J. (2024). Feed the Future Innovation Lab for Genomics to Improve Poultry: a holistic approach to improve indigenous chicken production focusing on resilience to Newcastle disease. *World's Poultry Science Journal*, 80(2), 273-297. [CrossRef]
- [133] Smith, J. (2019). Genomics of avian viral infections. *Genes*, 10(10), 814. [CrossRef]
- [134] Feng, A., Akter, S., Leigh, S. A., Wang, H., Pharr, G. T., Evans, J., ... & Wan, X. F. (2023). Genomic diversity, pathogenicity and antimicrobial resistance of *Escherichia coli* isolated from poultry in the southern United States. *BMC microbiology*, 23(1), 15. [CrossRef]
- [135] Olaniyan, O. F., Oladejo, O. A., Adeola, A. C., Bello, S. F., Bashiru, H. A., & Oseni, S. O. (2024). Integration of genomics into community-based breeding programmes for chickens: an overview of opportunities, challenges, and potential benefits. *World's Poultry Science Journal*, 80(3), 977-997. [CrossRef]
- [136] Wadood, A. A., & Zhang, X. (2024). The Omics Revolution in Understanding Chicken Reproduction: A Comprehensive Review. *Current Issues in Molecular Biology*, 46(6), 6248-6266. [CrossRef]
- [137] Long, J. A. (2020). The 'omics' revolution: Use of genomic, transcriptomic, proteomic and metabolomic tools to predict male reproductive traits that impact fertility in livestock and poultry. *Animal Reproduction Science*, 220, 106354. [CrossRef]
- [138] Nematbakhsh, S., Pei, C. P., Nordin, N., Selamat, J., Idris, L. H., & Razis, A. F. A. (2024). Identification and validation of novel breed-specific biomarker for the purpose of village chicken authentication using genomics approaches. *Poultry Science*, 103(11), 104128. [CrossRef]
- [139] Barros, C. P., Derks, M. F., Mohr, J., Wood, B. J., Crooijmans, R. P., Megens, H. J., ... & Groenen, M. A. (2023). A new haplotype-resolved turkey genome to enable turkey genetics and genomics research. *GigaScience*, 12, giad051. [CrossRef]
- [140] Li, M., Sun, C., Xu, N., Bian, P., Tian, X., Wang, X., ... & Yang, N. (2022). De novo assembly of 20 chicken genomes reveals the undetectable phenomenon for thousands of core genes on microchromosomes and subtelomeric regions. *Molecular Biology and Evolution*, 39(4), msac066. [CrossRef]
- [141] Che, T., Li, J., Li, X., Wang, Z., Zhang, X., Yang, W., ... & Zhang, W. (2024). Haplotype-resolved assembly of the mule duck genome using high-fidelity sequencing technology. *Plos one*, 19(7), e0305914. [CrossRef]
- [142] Guo, X., He, X. X., Chen, H., Wang, Z. C., Li, H. F., Wang, J. X., ... & Jiang, R. S. (2021). Revisiting the evolutionary history of domestic and wild ducks based on genomic analyses. *Zoological Research*, 42(1), 43. [CrossRef]
- [143] Chang, G., Yuan, X., Guo, Q., Bai, H., Cao, X., Liu, M., ... & Chen, G. (2023). The first crested duck genome reveals clues to genetic compensation and crest cushion formation. *Genomics, Proteomics & Bioinformatics*, 21(3), 483-500. [CrossRef]
- [144] Lin, R., Li, H., Lin, W., Yang, F., Bao, X., Pan, C., ... & Lin, W. (2024). Whole-genome selection signature differences between Chaohu and Ji'an red ducks. *BMC genomics*, 25(1), 522. [CrossRef]
- [145] Du, X., Lai, S., Zhao, W., Xu, X., Xu, W., Zeng, T., ... & Lu, L. (2022). Single-cell RNA sequencing revealed the liver heterogeneity between egg-laying duck and ceased-laying duck. *BMC genomics*, 23(1), 857. [CrossRef]
- [146] Zhang, X., Zhu, T., Wang, L., Lv, X., Yang, W., Qu, C., ... & Qu, L. (2023). Genome-wide association study reveals the genetic basis of duck plumage colors. *Genes*, 14(4), 856. [CrossRef]
- [147] Yu, S., Liu, Z., Li, M., Zhou, D., Hua, P., Cheng, H., ... & Zhou, Z. (2023). Resequencing of a Pekin duck breeding population provides insights into the genomic response to short-term artificial selection. *Gigascience*, 12, giad016. [CrossRef]
- [148] Wen, J., Li, H., Wang, H., Yu, J., Zhu, T., Zhang, J., ... & Qu, L. (2023). Origins, timing and introgression of domestic geese revealed by whole genome data. *Journal of animal science and biotechnology*, 14(1), 26. [CrossRef]
- [149] Zhang, Y., Ni, H., Xie, H., Yin, Y., Zheng, J., Dong, L., ... & Sun, X. (2022). De novo assembly of a wild swan goose (*Anser cygnoides*) genome. *Animal Genetics*, 53(6), 878-880. [CrossRef]
- [150] Zhou, Y., Mabrouk, I., Ma, J., Liu, Q., Song, Y.,

- Xue, G., ... & Sun, Y. (2024). Chromosome-level genome sequencing and multi-omics of the Hungarian White Goose (*Anser anser domesticus*) reveals novel miRNA-mRNA regulation mechanism of waterfowl feather follicle development. *Poultry Science*, 103(9), 103933. [CrossRef]
- [151] Liu, J., Xiao, Y., Ren, P., Zhang, S., Liu, Y., & Zhu, M. (2023). Integrating genomics and transcriptomics to identify candidate genes for high egg production in Wulong geese (*Anser cygnoides orientalis*). *BMC genomics*, 24(1), 481. [CrossRef]
- [152] Zhang, L., Li, H., Tang, B., Zhao, X., Wu, Y., Jiang, T., ... & Wang, L. (2023). Genomic signatures reveal selection in Chinese and European domesticated geese. *Animal Genetics*, 54(6), 763-771. [CrossRef]
- [153] Guinan, F. L., Wiggans, G. R., Norman, H. D., Dürr, J. W., Cole, J. B., Van Tassell, C. P., ... & Lourenco, D. (2023). Changes in genetic trends in US dairy cattle since the implementation of genomic selection. *Journal of dairy science*, 106(2), 1110-1129. [CrossRef]
- [154] Kertz, N. C., Banerjee, P., Dyce, P. W., & Diniz, W. J. (2023). Harnessing genomics and transcriptomics approaches to improve female fertility in beef cattle—a review. *Animals*, 13(20), 3284. [CrossRef]
- [155] Du, X., Sun, Y., Fu, T., Gao, T., & Zhang, T. (2024). Research Progress and Applications of Bovine Genome in the Tribe Bovini. *Genes*, 15(4), 509. [CrossRef]
- [156] Tian, R., Mahmoodi, M., Tian, J., Esmailizadeh Koshkoiyeh, S., Zhao, M., Saminzadeh, M., ... & Esmailizadeh, A. (2024). Leveraging Functional Genomics for Understanding Beef Quality Complexities and Breeding Beef Cattle for Improved Meat Quality. *Genes*, 15(8), 1104. [CrossRef]
- [157] Zhao, J., Wang, C., Hu, J., Ma, R., Yu, B., Zhao, W., ... & Zhang, J. (2024). Integrated metagenomics and metabolomics analyses revealed biomarkers in β -casein A2A2-type cows. *Frontiers in Veterinary Science*, 11, 1438717. [CrossRef]
- [158] Billa, P. A., Faulconnier, Y., Ye, T., Bourdon, C., Pires, J. A., & Leroux, C. (2021). Nutrigenomic analyses reveal miRNAs and mRNAs affected by feed restriction in the mammary gland of midlactation dairy cows. *PLoS One*, 16(4), e0248680. [CrossRef]
- [159] Zavadilová, L., Kašná, E., Krupová, Z., & Klímová, A. (2021). Health traits in current dairy cattle breeding: A review. *Czech Journal of Animal Science*, 66(7), 235-250. [CrossRef]
- [160] Yin, S., Li, Z., Yang, F., Guo, H., Zhao, Q., Zhang, Y., ... & He, J. (2023). A comprehensive genomic analysis of Chinese indigenous ningxiang pigs: genomic breed compositions, runs of homozygosity, and beyond. *International Journal of Molecular Sciences*, 24(19), 14550. [CrossRef]
- [161] Li, G., Liu, Y., Feng, X., Diao, S., Zhong, Z., Li, B., ... & Zhang, Z. (2024). Integrating Multiple Database Resources to Elucidate the Gene Flow in Southeast Asian Pig Populations. *International Journal of Molecular Sciences*, 25(11), 5689. [CrossRef]
- [162] Piórkowska, K., & Ropka-Molik, K. (2024). Trends and Prospects in Pig Genomics and Genetics. *Genes*, 15(10), 1292. [CrossRef]
- [163] Johnsson, M., Hickey, J. M., & Jungnickel, M. K. (2024). Building in vitro tools for livestock genomics: chromosomal variation within the PK15 cell line. *BMC genomics*, 25(1), 49. [CrossRef]
- [164] Flack, N., Hughes, L., Cassens, J., Enriquez, M., Gebeyehu, S., Alshagawi, M., ... & Faulk, C. (2024). The genome of Przewalski's horse (*Equus ferus przewalskii*). *G3: Genes, Genomes, Genetics*, 14(8), jkae113. [CrossRef]
- [165] Littiere, T. O., Castro, G. H., Rodriguez, M. D. P. R., Bonafé, C. M., Magalhães, A. F., Faleiros, R. R., ... & Verardo, L. L. (2020). Identification and functional annotation of genes related to horses' performance: from GWAS to post-GWAS. *Animals*, 10(7), 1173. [CrossRef]
- [166] Polak, G., Gurgul, A., Jasielczuk, I., Szmatoła, T., Krupiński, J., & Bugno-Poniewierska, M. (2021). Suitability of pedigree information and genomic methods for analyzing inbreeding of Polish cold-blooded horses covered by conservation programs. *Genes*, 12(3), 429. [CrossRef]
- [167] Fröhlich, D. E., Wallner, B., Juras, R., Cothran, E. G., & Velie, B. D. (2024). Relatedness and genomic inbreeding in a sample of Timor ponies. *Journal of Equine Veterinary Science*, 133, 105016. [CrossRef]
- [168] Saleem, A., Khan, A. A., Dar, M. A., Ganaie, A. M., Beigh, Y. A., & Ahmad, S. M. (2022). Nutrigenomics in livestock sector and its human-animal interface—a review. *Vet. Anim. Sci*, 17, 100262.
- [169] Taylor, J. F., Schnabel, R. D., & Sutovsky, P. (2018). Genomics of bull fertility. *animal*, 12(s1), s172-s183. [CrossRef]
- [170] Zhang, F., Shi, C., He, Q., Zhu, L., Zhao, J., Yao, W., ... & Luo, J. (2024). Integrated analysis of genomics and transcriptomics revealed the genetic basis for goaty flavor formation in goat milk. *Genomics*, 116(4), 110873. [CrossRef]
- [171] Li, C., Wu, Y., Chen, B., Cai, Y., Guo, J., Leonard, A. S., ... & Wang, X. (2022). Markhor-derived introgression of a genomic region encompassing PAPSS2 confers high-altitude adaptability in Tibetan goats. *Molecular Biology and Evolution*, 39(12), msac253. [CrossRef]
- [172] VarGoats Consortium, Nijman, I. J., Rosen, B. D., Bardou, P., Faraut, T., Cumer, T., ... & Anton, I. (2022). Geographical contrasts of Y-chromosomal haplogroups from wild and domestic goats reveal ancient migrations and recent introgressions. *Molecular ecology*, 31(16), 4364-4380. [CrossRef]
- [173] Negro, A., Cesarani, A., Cortellari, M., Bionda, A., Fresi, P., Macciotta, N. P. P., ... & Crepaldi, P. (2024). A

- comparison of genetic and genomic breeding values in Saanen and Alpine goats. *animal*, 18(4), 101118. [CrossRef]
- [174] Denoyelle, L., Talouarn, E., Bardou, P., Colli, L., Alberti, A., Danchin, C., ... & Tosser-Klopp, G. (2021). VarGoats project: a dataset of 1159 whole-genome sequences to dissect *Capra hircus* global diversity. *Genetics Selection Evolution*, 53(1), 86. [CrossRef]
- [175] Lyons, A., Brown, J., & Davenport, K. M. (2024). Single-cell sequencing technology in ruminant livestock: challenges and opportunities. *Current Issues in Molecular Biology*, 46(6), 5291-5306. [CrossRef]
- [176] Park, J. S., Lee, K. Y., & Han, J. Y. (2020). Precise genome editing in poultry and its application to industries. *Genes*, 11(10), 1182. [CrossRef]
- [177] Alvarenga, M., D'Elia, A. K. P., Rocha, G., Arantes, C. A., Henning, F., de Vasconcelos, A. T. R., & Solé-Cava, A. M. (2024). Mitochondrial genome structure and composition in 70 fishes: a key resource for fisheries management in the South Atlantic. *BMC genomics*, 25(1), 215. [CrossRef]
- [178] Papa, Y., Wellenreuther, M., Morrison, M. A., & Ritchie, P. A. (2023). Genome assembly and isoform analysis of a highly heterozygous New Zealand fisheries species, the tarakihi (*Nemadactylus macropterus*). *G3*, 13(2), jkac315. [CrossRef]
- [179] Bertram, A., Fairclough, D., Sandoval-Castillo, J., Brauer, C., Fowler, A., Wellenreuther, M., & Beheregaray, L. B. (2022). Fisheries genomics of snapper (*Chrysophrys auratus*) along the west Australian coast. *Evolutionary Applications*, 15(7), 1099-1114. [CrossRef]
- [180] Blommaert, J., Sandoval-Castillo, J., Beheregaray, L. B., & Wellenreuther, M. (2024). Peering into the gaps: Long-read sequencing illuminates structural variants and genomic evolution in the Australasian snapper. *Genomics*, 116(5), 110929. [CrossRef]
- [181] Timm, L. E., Jackson, T. L., Browder, J. A., & Bracken-Grissom, H. D. (2021). Population Genomics of the Commercially Important Gulf of Mexico Pink Shrimp *Farfantepenaeus duorarum* Support Models of Juvenile Transport Around the Florida Peninsula. *Frontiers in Ecology and Evolution*, 9, 659134. [CrossRef]
- [182] Ferrette, B. L. D. S., Coimbra, R. T., Winter, S., De Jong, M. J., Williams, S. M., Coelho, R., ... & Janke, A. (2023). Seascape Genomics and Phylogeography of the Sailfish (*Istiophorus platypterus*). *Genome Biology and Evolution*, 15(4), evad042. [CrossRef]
- [183] Woodings, L. N., Murphy, N. P., Liggins, G. W., Miller, M. E., Ballinger, G. M., Lau, S. C., & Strugnelli, J. M. (2021). Population genomics of the Eastern Rock Lobster, *Sagmariasus verreauxi*, during spawning stock recovery from over-exploitation. *ICES Journal of Marine Science*, 78(7), 2448-2459. [CrossRef]
- [184] Timm, L. E., Tucker, N., Rix, A., LaBua, S., López, J. A., Boswell, K. M., & Glass, J. R. (2023). The untapped potential of seascape genomics in the North Pacific. *Frontiers in Conservation Science*, 4, 1249551. [CrossRef]
- [185] Van Oppen, M. J., & Coleman, M. A. (2022). Advancing the protection of marine life through genomics. *PLoS Biology*, 20(10), e3001801. [CrossRef]
- [186] Caccavo, J. A., Arantes, L. S., Celemín, E., Mbedi, S., Sparmann, S., & Mazzoni, C. J. (2024). Whole-genome resequencing improves the utility of otoliths as a critical source of DNA for fish stock research and monitoring. *Molecular Ecology Resources*, 24(8), e14013. [CrossRef]
- [187] Casas, L., & Saborido-Rey, F. (2023). A review of genomics methods and bioinformatics tools for the analysis of close-kin mark-recapture. *Frontiers in Marine Science*, 10, 1113870. [CrossRef]
- [188] Casas, L., Hanel, R., Piferrer, F., & Saborido-Rey, F. (2023). Prospects and challenges for the implementation of HTS genetic methods in fisheries research surveys and stock assessments. *Frontiers in Marine Science*, 10, 1238133. [CrossRef]
- [189] Andersson, L., Bekkevold, D., Berg, F., Farrell, E. D., Felkel, S., Ferreira, M. S., ... & Pettersson, M. (2024). How fish population genomics can promote sustainable fisheries: A road map. *Annual Review of Animal Biosciences*, 12(1), 1-20. [CrossRef]
- [190] Attard, C. R., Sandoval-Castillo, J., Gilligan, D. M., Unmack, P. J., Faulks, L. K., & Beheregaray, L. B. (2022). Genomics outperforms genetics to manage mistakes in fisheries stocking of threatened species. *Biodiversity and Conservation*, 31(3), 895-908. [CrossRef]
- [191] Cao, H. X., Vu, G. T. H., & Gailing, O. (2022). From genome sequencing to CRISPR-based genome editing for climate-resilient forest trees. *International Journal of Molecular Sciences*, 23(2), 966. [CrossRef]
- [192] Sharma, U., Sankhyani, H. P., Kumari, A., Thakur, S., Thakur, L., Mehta, D., ... & Sankhyani, N. (2024). Genomic selection: A revolutionary approach for forest tree improvement in the wake of climate change. *Euphytica*, 220(1), 9. [CrossRef]
- [193] Mostert-O'Neill, M. M., Tate, H., Reynolds, S. M., Mphahlele, M. M., van den Berg, G., Verry, S. D., ... & Myburg, A. A. (2022). Genomic consequences of artificial selection during early domestication of a wood fibre crop. *New Phytologist*, 235(5), 1944-1956. [CrossRef]
- [194] Anders, C., Hoengenaert, L., & Boerjan, W. (2023). Accelerating wood domestication in forest trees through genome editing: Advances and prospects. *Current opinion in plant biology*, 71, 102329. [CrossRef]
- [195] Ahmar, S., Ballesta, P., Ali, M., & Mora-Poblete, F. (2021). Achievements and challenges of genomics-assisted breeding in forest trees: From marker-assisted selection to genome editing. *International Journal of Molecular Sciences*, 22(19), 10583. [CrossRef]
- [196] Burley, J. T., Kellner, J. R., Hubbell, S. P., & Faircloth,

- B. C. (2021). Genome assemblies for two Neotropical trees: *Jacaranda copaia* and *Handroanthus guayacan*. *G3*, 11(2), jkab010. [[CrossRef](#)]
- [197] Niu, S., & Li, Q. (2024). From haploid to reference: a new milestone in poplar genomics. *Forestry Research*, 4, e020. [[CrossRef](#)]
- [198] Tumas, H., Iliska, J. J., Gérardi, S., Laroche, J., A'Hara, S., Boyle, B., ... & MacKay, J. J. (2024). High-density genetic linkage mapping in Sitka spruce advances the integration of genomic resources in conifers. *G3: Genes, Genomes, Genetics*, 14(4), jkae020. [[CrossRef](#)]
- [199] Capblancq, T., Sękiewicz, K., & Dering, M. (2024). Forest genomics in the Caucasus through the lens of its dominant tree species—*Fagus orientalis*. *Molecular Ecology*, 33(16), e17475. [[CrossRef](#)]
- [200] Luo, L., Ye, P., Lin, Q., Liu, M., Hao, G., Wei, T., & Sahu, S. K. (2024). From sequences to sustainability: exploring dipterocarp genomes for oleoresin production, timber quality, and conservation. *Plant Science*, 112139. [[CrossRef](#)]
- [201] Han, B., Wang, L., Xian, Y., Xie, X. M., Li, W. Q., Zhao, Y., ... & Jia, K. H. (2022). A chromosome-level genome assembly of the Chinese cork oak (*Quercus variabilis*). *Frontiers in Plant Science*, 13, 1001583. [[CrossRef](#)]
- [202] Matallana-Ramirez, L. P., Whetten, R. W., Sanchez, G. M., & Payn, K. G. (2021). Breeding for climate change resilience: a case study of loblolly pine (*Pinus taeda* L.) in North America. *Frontiers in plant science*, 12, 606908. [[CrossRef](#)]
- [203] Shirasawa, K., Mishima, K., Hirakawa, H., Hirao, T., Tsubomura, M., Nagano, S., ... & Takahashi, M. (2024). Haplotype-resolved de novo genome assemblies of four coniferous tree species. *Journal of Forest Research*, 29(2), 151-157. [[CrossRef](#)]
- [204] Feng, J., Dan, X., Cui, Y., Gong, Y., Peng, M., Sang, Y., ... & Wang, J. (2024). Integrating evolutionary genomics of forest trees to inform future tree breeding amidst rapid climate change. *Plant Communications*, 5(10), 101044. [[CrossRef](#)]
- [205] Abiri, R., Abdul-Hamid, H., & Leung, D. W. (2023). Advanced insights into environmental stress and tolerance mechanisms in forest trees. *Frontiers in Plant Science*, 14, 1273663. [[CrossRef](#)]



Yufei Yang focuses on genomics, gene editing, agricultural breeding and their interconnections. (Email: 2303020124@st.btbu.edu.cn)